

Note: In this problem set, expressions in green cells match corresponding expressions in the text answers.

2 - 13 Verification of Solutions

Verify (by substitution) that the given function is a solution of the PDE. Sketch or graph the solution as a surface in space.

2 - 5 Wave Equation (1) with suitable c

$$3. u = \cos 4t \sin 2x$$

```
Clear["Global`*"]
```

```
u[x_, t_] = Cos[4 t] Sin[2 x]
```

```
Cos[4 t] Sin[2 x]
```

```
d1 = D[u[x, t], {t, 2}]
```

```
-16 Cos[4 t] Sin[2 x]
```

```
d2 = D[u[x, t], {x, 2}]
```

```
-4 Cos[4 t] Sin[2 x]
```


```
d1 == c^2 d2 (* 1D wave equation *)
```

```
-16 Cos[4 t] Sin[2 x] == -4 c^2 Cos[4 t] Sin[2 x]
```

```
Solve[-16 Cos[4 t] Sin[2 x] == -4 c^2 Cos[4 t] Sin[2 x], {c}]
```

```
{{c -> -2}, {c -> 2}}
```

```
Plot3D[Cos[4 t] Sin[2 x], {x, 0, Pi}, {t, 0, Pi}]
```



The value of the constant, c , is the key to the description of the particular solution u .

$$5. u = \sin at \sin bx$$

```
Clear["Global`*"]
```

```
u[x_, t_] = Sin[a t] Sin[b x]
Sin[a t] Sin[b x]
```

```
d1 = D[u[x, t], {t, 2}]
-a^2 Sin[a t] Sin[b x]
```

```
d2 = D[u[x, t], {x, 2}]
-b^2 Sin[a t] Sin[b x]
```


```
d1 == c^2 d2 (* 1D wave equation *)
-a^2 Sin[a t] Sin[b x] == -b^2 c^2 Sin[a t] Sin[b x]
```

```
Solve[-a^2 Sin[a t] Sin[b x] == -b^2 c^2 Sin[a t] Sin[b x], {c}]
```

```
{{c -> -a/b}, {c -> a/b}}
```

```
subeq = u[x, t] /. {a -> 2, b -> 3}
Sin[2 t] Sin[3 x]
```

```
Plot3D[subeq, {x, 0, Pi}, {t, 0, Pi}]
```



6 - 9 Heat Equation (2) with suitable c

$$7. u = e^{-\omega^2 c^2 t} \sin x$$

```
Clear["Global`*"]
```

```
u[x_, t_] = e^{-\omega^2 c^2 t} Sin[x]
e^{-c^2 t \omega^2} Sin[x]
```

```
d1 = D[u[x, t], {t}]
-c^2 e^{-c^2 t \omega^2} \omega^2 Sin[x]
```


```
d2 = D[u[x, t], {x, 2}]
-e^{-c^2 t \omega^2} Sin[x]
```

```
d1 == c^2 d2 (* 1D heat equation *)
-c^2 e^{-c^2 t \omega^2} \omega^2 Sin[x] == -c^2 e^{-c^2 t \omega^2} Sin[x]
```

I can't get Solve to give me what I want here. By inspection, c can take on any value, with $\omega = 1$ or -1 .

```
subeq = u[x, t] /. {c -> 2, \omega -> 1}
e^{-4 t} Sin[x]
```

```
Plot3D[subeq, {x, 0, Pi}, {t, 0, Pi}]
```



9. $u = e^{-\pi^2 t} \cos 25x$

```
Clear["Global`*"]
```

```
u[x_, t_] = e^{-\pi^2 t} Cos[25 x]
e^{-\pi^2 t} Cos[25 x]
```

```
d1 = D[u[x, t], {t}]
-e^{-\pi^2 t} \pi^2 Cos[25 x]
```

```
d2 = D[u[x, t], {x, 2}]
-625 e^{-\pi^2 t} Cos[25 x]
```

```
d1 == c^2 d2 (* 1D heat equation *)
-e^{-\pi^2 t} \pi^2 Cos[25 x] == -625 c^2 e^{-\pi^2 t} Cos[25 x]
```


```
Solve[-e^{-\pi^2 t} \pi^2 Cos[25 x] == -625 c^2 e^{-\pi^2 t} Cos[25 x], {c}]
```

```
{ {c -> -\frac{\pi}{25}}, {c -> \frac{\pi}{25}} }
```

One value of c agrees with the text answer. Mathematica adds the negative value, perhaps overlooked by the text.

```
subeq = u[x, t] /. {c -> \pi / 25}
e^{-\pi^2 t} Cos[25 x]
```

```
Plot3D[subeq, {x, 0, Pi}, {t, 0, Pi}]
```



10 - 13 Laplace Equation (3)

10. $u = e^x \cos y, e^x \sin y$

```
Clear["Global`*"]
```

```
ucos[x_, y_] = ex Cos[y]
```

```
ex Cos[y]
```

```
d1 = D[ucos[x, y], {x, 2}]
```

```
ex Cos[y]
```

```
d2 = D[ucos[x, y], {y, 2}]
```

```
-ex Cos[y]
```

```
eqc = d1 + d2 (* 2D Laplace equation *)
```

```
0
```

```
usin[x_, y_] = ex Sin[y]
```

```
ex Sin[y]
```

```
d3 = D[usin[x, y], {x, 2}]
```

```
ex Sin[y]
```


```
d4 = D[usin[x, y], {y, 2}]
```

```
-ex Sin[y]
```

```
eqs = d3 + d4 (* 2D Laplace equation *)
```

```
0
```

```
Plot3D[{ucos[x, y], usin[x, y]}, {x, 0, Pi}, {y, 0, Pi}]
```



I wasn't supposed to work this even-numbered problem, but in view of difficulties encountered in the next one, I'll leave this one in for now.

11. $u = \arctan(y/x)$

```
Clear["Global`*"]
```

```
u[x_, y_] = ArcTan[y / x]
```

```
ArcTan[ $\frac{y}{x}$ ]
```

```
d1 = D[u[x, y], {x, 2}]
```

$$-\frac{2y^3}{x^5 \left(1 + \frac{y^2}{x^2}\right)^2} + \frac{2y}{x^3 \left(1 + \frac{y^2}{x^2}\right)}$$

```
d2 = D[u[x, y], {y, 2}]
```


$$-\frac{2y}{x^3 \left(1 + \frac{y^2}{x^2}\right)^2}$$

```
eq2 = d1 + d2 == 0 (* 2D Laplace equation *)
```

$$-\frac{2y}{x^3 \left(1 + \frac{y^2}{x^2}\right)^2} - \frac{2y^3}{x^5 \left(1 + \frac{y^2}{x^2}\right)^2} + \frac{2y}{x^3 \left(1 + \frac{y^2}{x^2}\right)} == 0$$


An answer to this problem is omitted in the text. I can try to plot it (the Laplace surface).

```
Plot3D[- $\frac{2 y}{x^3 \left(1 + \frac{y^2}{x^2}\right)^2}$  -  $\frac{2 y^3}{x^5 \left(1 + \frac{y^2}{x^2}\right)^2}$  +  $\frac{2 y}{x^3 \left(1 + \frac{y^2}{x^2}\right)}$ , {x, 0, Pi}, {y, 0, Pi}]
```



The one that was supposed to be plotted is the solution, i.e. the given function:

```
Plot3D[ArcTan[y/x], {x, -1, 1}, {y, -1, 1}]
```



13. $u = x/(x^2 + y^2)$, $y/(x^2 + y^2)$

```
Clear["Global`*"]
```

```
ux[x_, y_] = x / (x^2 + y^2)
```

$$\frac{x}{x^2 + y^2}$$

```
d1 = D[ux[x, y], {x, 2}]
```

$$-\frac{4x}{(x^2 + y^2)^2} + x \left(\frac{8x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right)$$

```
d2 = D[ux[x, y], {y, 2}]
```

$$x \left(\frac{8y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right)$$

```
eq2 = d1 + d2 == 0 (* 2D Laplace equation, the sum = 0 *)
```

$$-\frac{4x}{(x^2 + y^2)^2} + x \left(\frac{8x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) + x \left(\frac{8y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) == 0$$

$$uy[x_, y_] = y / (x^2 + y^2)$$

$$\frac{y}{x^2 + y^2}$$

$$d3 = D[uy[x, y], {x, 2}]$$

$$y \left(\frac{8 x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right)$$

$$d4 = D[uy[x, y], {y, 2}]$$

$$- \frac{4 y}{(x^2 + y^2)^2} + y \left(\frac{8 y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right)$$

$$eq3 = d3 + d4 == 0 \quad (* \text{ 2D Laplace equation, the sum} = 0 *)$$

$$- \frac{4 y}{(x^2 + y^2)^2} + y \left(\frac{8 x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) + y \left(\frac{8 y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) == 0$$

To have a look at the surfaces that makes the Laplace equation true:

$$\text{plot1} = \text{Plot3D}[$$

$$- \frac{4 x}{(x^2 + y^2)^2} + x \left(\frac{8 x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) + x \left(\frac{8 y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) == 0,$$


$$\{x, -1, 1\}, \{y, -1, 1\}];$$

$$\text{plot2} = \text{Plot3D}[$$

$$- \frac{4 y}{(x^2 + y^2)^2} + y \left(\frac{8 x^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) + y \left(\frac{8 y^2}{(x^2 + y^2)^3} - \frac{2}{(x^2 + y^2)^2} \right) == 0,$$

$$\{x, -1, 1\}, \{y, -1, 1\}];$$

Show[plot1] Show[plot2]




$$\text{plot3} = \text{Plot3D}\left[\frac{x}{x^2 + y^2}, \{x, -1, 1\}, \{y, -1, 1\}\right];$$

$$\text{plot4} = \text{Plot3D}\left[\frac{y}{x^2 + y^2}, \{x, -1, 1\}, \{y, -1, 1\}\right];$$

Show[plot3] Show[plot4]

And at the given functions:



No answer to this problem appears in the text's answer appendix.

15. Boundary value problem

Verify that the function $u(x,y) = a \log(x^2 + y^2) + b$ satisfies Laplace's equation (3) and determine a and b so that u satisfies the boundary conditions $u=110$ on the circle $x^2 + y^2=100$.

```
Clear["Global`*"]
```

This one is worked in the s.m.

$$u[x_, y_] = a \text{Log}[x^2 + y^2] + b$$

$$b + a \text{Log}[x^2 + y^2]$$

$$d1 = D[u[x, y], \{x, 2\}]$$

$$a \left(-\frac{4x^2}{(x^2 + y^2)^2} + \frac{2}{x^2 + y^2} \right)$$

$$d2 = D[u[x, y], \{y, 2\}]$$

$$a \left(-\frac{4y^2}{(x^2 + y^2)^2} + \frac{2}{x^2 + y^2} \right)$$

```
FullSimplify[d1 + d2]
```

0

The Laplace equation equality is verified. The function u is a solution. Now for the boundary values.

```
Solve[a Log[100] + b == 110, {b}]
```

{b -> 110 - a Log[100]}


```
Solve[a Log[100] + b == 110, {a}]
```

$$\left\{ \left\{ a \rightarrow \frac{110 - b}{\text{Log}[100]} \right\} \right\}$$

I was not overly pleased with the way the discovery of the constants a and b needed to be done. I could not find a way to do it in one step.

16 - 23 PDEs Solvable as ODEs

This happens if a PDE involves derivatives with respect to one variable only (or can be transformed to such a form), so that the other variable(s) can be treated as parameter(s).

Solve for $u = u(x,y)$:

$$17. u_{xx} + 16\pi^2 u = 0$$

```
Clear["Global`*"]
```

```
eqn = D[u[x, y], {x, 2}] + 16 π2 u[x, y] == 0
```

```
16 π2 u[x, y] + u(2,0)[x, y] == 0
```

```
sol = DSolve[eqn, u[x, y], {x, y}]
```

$$\left\{ \left\{ u[x, y] \rightarrow \text{Cos}[4\pi x] C[1][y] + \text{Sin}[4\pi x] C[2][y] \right\} \right\}$$

Even though the independent variable y does not make an active appearance, its presence must be directly acknowledged in order to get its representation shown in the solution. The answer matches the text's.

$$19. u_y + y^2 u = 0$$

```
Clear["Global`*"]
```

```
eqn = D[u[x, y], {y}] + y2 u[x, y] == 0
```

```
y2 u[x, y] + u(0,1)[x, y] == 0
```

```
sol = DSolve[eqn, u[x, y], {x, y}]
```

$$\left\{ \left\{ u[x, y] \rightarrow e^{-\frac{y^3}{3}} C[1][x] \right\} \right\}$$

The above answer matches the text's.

$$21. u_{yy} + 6u_y + 13u = 4e^{3y}$$

```
Clear["Global`*"]
```

```
eqn = D[u[x, y], {y, 2}] + 6 D[u[x, y], {y}] + 13 u[x, y] - 4 e3y == 0
```

```
-4 e3y + 13 u[x, y] + 6 u(0,1)[x, y] + u(0,2)[x, y] == 0
```

```
sol = Simplify[DSolve[eqn, u[x, y], {x, y}]]
```

$$\left\{ \left\{ u[x, y] \rightarrow \frac{1}{10} e^{-3y} \left(e^{6y} + 10 \sin[2y] C[1][x] + 10 \cos[2y] C[2][x] \right) \right\} \right\}$$

The above answer matches the text's.

$$23. x^2 u_{xx} + 2x u_x - 2u = 0$$

```
Clear["Global`*"]
```

```
eqn = x^2 D[u[x, y], {x, 2}] + 2 x D[u[x, y], {x}] - 2 u[x, y] == 0
```

```
- 2 u[x, y] + 2 x u^{(1,0)}[x, y] + x^2 u^{(2,0)}[x, y] == 0
```

```
sol = Simplify[DSolve[eqn, u[x, y], {x, y}]]
```

$$\left\{ \left\{ u[x, y] \rightarrow x C[1][y] + \frac{C[2][y]}{x^2} \right\} \right\}$$

The above answer matches the text's.

25. System of PDEs

$$\text{Solve } u_{xx} = 0, u_{yy} = 0$$

```
Clear["Global`*"]
```

```
eqn1 = D[u[x, y], {x, 2}] == 0
```

```
u^{(2,0)}[x, y] == 0
```

```
eqn2 = D[u[x, y], {y, 2}] == 0
```

```
u^{(0,2)}[x, y] == 0
```

```
DSolve[{{u^{(2,0)}[x, y] == 0}, {u^{(0,2)}[x, y] == 0}}, u[x, y], {x, y}]
```

```
DSolve[{{u^{(2,0)}[x, y] == 0}, {u^{(0,2)}[x, y] == 0}}, u[x, y], {x, y}]
```

After trying several variations in formatting, I find that Mathematica 10 will not do this differential equation system. I find that Mathematica 11 won't do it either, and neither will WolframAlpha.

```
h1 = DSolve[eqn1, u[x, y], {x, y}]
```

```
{{u[x, y] → C[1][y] + x C[2][y]}}
```

```
h2 = DSolve[eqn2, u[x, y], {x, y}]
```

```
{{u[x, y] → C[1][x] + y C[2][x]}}
```

$$\text{tot} = C[1][y] + x C[2][y] + C[3][x] + y C[4][x]$$

The simplicity of the system allows it to be done by hand, by adding the partial solutions. In the above yellow cell the $C[2]$ and $C[4]$ terms need to be combined, and there is no isolated arbitrary constant. With these modifications, it would match the text answer.